Biologic Therapy



6308 Benjamin Rd #709, Tampa, FL 33634 Phone: (727) 896-0001

Fax: (727) 896-0002

Patient Information		Prescriber + Shipping Information	
Patient name: DOB:		Prescriber name:	
Sex: Female Male SSN: Ethnicity:		NPI:	
Language: Wt: kg			
Address:		Apt/Suite: City: State: _	
Apt/Suite: City: State: Zip:			
Phone:Alternate:		Contact:	
Caregiver name: Relation:		Phone: Alternate:	
Bin: PCN:		Fax:	
ID: Group #:		Email:	
Please fax a copy of front and back of th	e insurance card(s).	If shipping to prescriber: ☐ First Fill ☐ Always ☐ Ne	ever
Comorbidities:			
Concomitant Medications:			
Allergies: NKDA Other:			
Prescription	Form/Strength/Directions F	or use	Quantity Refills
∏Xifaxan	□ 200 mg mg po □ BID □ TID for days □ 550 mg mg po □ BID □ TID for days		
□Simponi	☐ Initial Dosing: 200 mg SC week 0 then 100mg SC week 2 ☐ Other: ☐ Ongoing treatment: 100mg SC every 4 weeks ☐ Other:		
☐ Cimzia (200mg Vials for PFS)	 □ Lyophilized vial □ Pre-Filled Syringe □ Initial Dosing: 400mg SC at 0, 2, 4 weeks (ongoing treatment below) □ Ongoing treatment (after initial dosing): 400mg every 4 weeks 		
□Stelara	□ 45mg PFS SC x1 followed by 45mg PFS SC in 4 weeks (For patients weighing < 100kg) □ 90mg PFS SC x1 followed by 90mg PFS SC in 4 weeks (For patients weighing > 100kg) Maintenance Dosing: □ 45mg PFS SC every 12 weeks □ 90mg PFS SC every 12 weeks □ 260mg □ 390mg □ 520mg followed by □ 90mg IV every 8 weeks		
☐ Humira (Supplied as 40mg pens or as PFS)	 Initial Dosing: Chron's/Ulcerative Colitis Starter Pack - six 40mg single dose pens 160mg (four 40mg injections) SC x one dose (day 1) then 80mg SC x one two weeks later (day 15) ☐ Ongoing treatment: ☐ Pen or ☐ PFS 40mg SC every other week (Start 29 days after initial dosing) ☐ Other: 		
□ Humira Pediatric (Supplied as PFS)	Initial dosing for children >/= 6 and >/= 40kg: □ 4x40 mg inj SC on Day 1 then 2x40mg inj SC 2 weeks later on Day 15 (tray of 6) □ 2x40 mg inj SC on Days 1 and 2 then 2x40mg inj SC 2 weeks later on Day 15 (tray of 6) Initial dosing for children >/= 6 and weighing 17kg to < 40kg: □ 2x40 mg inj SC on Day 1 then 1x40mg inj SC 2 weeks later on Day 15 (tray of 3) Maintenance: □ 40mg SC every other week □ 20mg SC every other week		
□Relistor	☐ 150mg tab ☐ 8mg/0.4ml Sol'n Sig:	, — ,	
☐ Other			
Information Needed to Obtain Prior Authorization			
Primary Diagnosis: ☐ ICD-10 ☐ Other:			
Weight:pounds Allergies:			
Failed Therapies: Please provide current list of medications:			
**For the form (tablets or capsules), unless otherwise specified, pharmacy preference/availiability (or insurance preference) will be dispensed.			
Per state-specific law, prescriptions will be dispensed as generic, if applicable, unless notated otherwise:			
Prescriber's Signature:			