

6308 Benjamin Rd #709, Tampa, FL 33634 Phone: (727) 896-0001 Fax: (727) 896-0002

Patient Information		Prescriber + Snipping information	
Patient name: DOB:		Prescriber name:	
Sex: Female Male SSN: Ethnicity:		NPI:	
Language: Wt:		Address:	
Address:		Apt/Suite: City: State:	
Address: City:	State: Zip:	I .	
Phone: Alternate:		Contact:	
Caregiver name: Relation:			
Bin: PCN:		Fax:	
ID: Group #:			
Please fax a copy of front and back of the insurance card(s). If shipping to prescriber: First Fill Always Never Comorbidities:			
Concomitant Medications:			
Allergies: NKDA Other:			
Prescription	Form/Strength/Directions F	For Use	Quantity Refills
∏Xifaxan	☐ 200 mg mg po ☐ BID	□ TID for days	
	550 mg mg po		
□Simponi	☐ Initial Dosing: 200 mg SC week 0 then 100mg SC week 2 ☐ Other: ☐ Ongoing treatment: 100mg SC every 4 weeks ☐ Other:		
□Cimzia	☐ Lyophilized vial ☐ Pre-Filled Syringe		
(200mg Vials for PFS)	☐ Initial Dosing: 400mg SC at 0, 2, 4 weeks (ongoing treatment below)		
	☐ Ongoing treatment (after initial dosing): 400mg every 4 weeks		
□Stelara	45mg PFS SC x1 followed by 45mg PFS SC in 4 weeks (For patients weighing < 100kg)		
	90mg PFS SC x1 followed by 90mg PFS SC in 4 weeks (For patients weighing > 100kg)		
	Maintenance Dosing: ☐ 45mg PFS SC every 12 weeks ☐ 90mg PFS SC every 12 weeks		
	☐ 260mg ☐ 390mg ☐ 520mg followed by ☐ 90mg IV every 8 weeks		
☐ Humira	☐ Initial Dosing: Chron's/Ulcerative Colitis Starter Pack - six 40mg single dose pens 160mg (four 40mg injections) SC x one dose (day 1) then 80mg SC x one two weeks later (day 15)		
(Supplied as 40mg pens or as PFS)			
	☐ Ongoing treatment: ☐ Pen or ☐ PFS		
	40mg SC every other week (Start 29 days after initial dosing)		
	☐ Other:		
☐Humira Pediatric	Initial dosing for children >/= 6 and >/= 40kg:		
(Supplied as PFS)	□ 4x40 mg inj SC on Day 1 then 2x40mg inj SC 2 weeks later on Day 15 (tray of 6) □ 2x40 mg inj SC on Days 1 and 2 then 2x40mg inj SC 2 weeks later on Day 15 (tray		
	of 6)		
	Initial dosing for children >/= 6 and weighing 17kg to < 40kg:		
	2x40 mg inj SC on Day 1 then 1x40mg inj SC 2 weeks later on Day 15 (tray of 3)		
	Maintenance: ☐ 40mg SC every ot	her week 🔲 20mg SC every other week	
Relistor	☐ 150mg tab ☐ 8mg/0.4ml Sol'n	for Inj 12mg/0.6ml Sol'n for Inj	
	Sig:		
Other			
Information Needed to Obtain Pr	ior Authorization		
Primary Diagnosis: ☐ ICD-10 ☐ Other:			
Weight:pounds Allergies:			
Failed Therapies: Please provide current list of medications:			
**For the form (tablets or capsules), unless otherwise specified, pharmacy preference/availiability (or insurance preference) will be dispensed.			
Per state-specific law, prescriptions will be dispensed as generic, if applicable, unless notated otherwise:			
Prescriber's Signature:Date:			
I authorize Wellgistics Pharmacy and its representatives to act as an agent to initiate and execute the insurance prior authorization and appeal process for this prescription and any future fills of the same prescription for the patient listed above. I understand that I can revoke this designation at any time by providing written notice to Wellgistics Pharmacy.			